



#6
RECEIVED
JUN 21 2002
TECH CENTER 1600
Sheet 1 of 3

Form PTO-1449 Modified

List of Patent and Publications Cited by Applicant
(Use several sheets if necessary)

U.S. Department of Commerce
Patent and Trademark Office

Client Matter No.
13259.00015

Serial No.
09/763,331

Applicant
Charles E. Martin and Andrew Mitchell

Filing Date
February 22, 2001

Group
TBA

U. S. PATENT DOCUMENTS

Examiner Initial		Document No.	Date	Name	Class	Subclass
RK	AA	5,057,419	10/1991	Martin et al.	435	134
RK	AB	5,380,831	1/1995	Adang et al.	536	23.71
RK	AC	5,500,365	3/1996	Fischoff et al.	435	240.4

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)


RK -	AD -	Dolphin, M.E., <i>et al.</i> , The translational signal database, TransTerm: more organisms, complete genomes. Nucleic Acids Research 25:246-247 (1997).
	AE -	Fox, B.G. <i>et al.</i> , Stearyl-acyl carrier protein Δ^9 desaturase from <i>Ricinus communis</i> is a diiron-oxo protein. Proc. Natl. Acad. Sci. USA 90:2486-2490 (1993).
	AF	Grayburn, W.S. <i>et al.</i> , Fatty Acid Alteration By A Δ^9 Desaturase In Transgenic Tobacco Tissue. Bio/Technology 10:675-678 (1992).
	AG	Hamada, T. <i>et al.</i> , Modification of fatty acid composition by over- and antisense-expression of a microsomal omega-3 fatty acid desaturase gene in transgenic tobacco. Transgenic Research 5:115-121 (1996) (abstract only)
	AH	Hamada, T. <i>et al.</i> , cDNA Cloning of a Wounding-Inducible Gene Encoding a Plastid ω -3 Fatty Acid Desaturase from Tobacco. Plant Cell Physiol. 37:606-611 (1996).
	AI	Hebsgaard, S.M. <i>et al.</i> , Splice site prediction in <i>Arabidopsis thaliana</i> pre-mRNA by combining local and global sequence information. Nucleic Acids Research 24:3439-3452 (1996).
	AJ	Heppard, E.P. <i>et al.</i> , Developmental and Growth Temperature Regulation of Two Different Microsomal ω -6 Desaturase Genes in Soybeans. Plant Physiol. 111:311-319 (1996).
	AK	Hugly, S. <i>et al.</i> , Enhanced Thermal Tolerance of Photosynthesis and Altered Chloroplast Ultrastructure in a Mutant of <i>Arabidopsis</i> Deficient in Lipid Desaturation. Plant Physiol. 90:1134-1142 (1989).
RK	AL	Kozak, M., Structural Features in Eukaryotic mRNAs That Modulate the Initiation of Translation. The Journal of Biological Chemistry 266:19867-19870 (1991).

EXAMINER

Russell Kalli

DATE CONSIDERED

1-8-03



RECEIVED
TECH CENTER 1600/2900
JUN 21 2002

Form PTO-1449 ModifiedClient Matter No.
13259.00015Serial No.
09/763,331List of Patent and Publications Cited by Applicant
(Use several sheets if necessary)Applicant
Charles E. Martin and Andrew MitchellU.S. Department of Commerce
Patent and Trademark OfficeFiling Date
February 22, 2001Group
TBA**OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)**


RK -	BA	McDonough, V.M. <i>et al.</i> , Specificity of Unsaturated Fatty Acid-regulated Expression of the <i>Saccharomyces cerevisiae</i> <i>OLE1</i> Gene. <i>The Journal of Biological Chemistry</i> 267:5931-5936 (1992).
g2	BB	Mitchell, A.G. and Martin, C.E., A Novel Cytochrome <i>b_s</i> -like Domain Is Linked to the Carboxyl Terminus of the <i>Saccharomyces cerevisiae</i> Δ-9 Fatty Acid Desaturase. <i>The Journal of Biological Chemistry</i> 270:29766-29771 (1995).
,	BC	Murata, N. and Wada, H., Acyl-lipid desaturases and their importance in the tolerance and acclimatization to cold of cyanobacteria. <i>Biochem J.</i> 308:1-8 (1995).
,	BD	Oh, C. <i>et al.</i> , <i>ELO2</i> and <i>ELO3</i> Homologues of the <i>Saccharomyces cerevisiae</i> <i>ELO1</i> Gene, Function in Fatty Acid Elongation and Are Required for Sphingolipid Formation. <i>The Journal of Biological Chemistry</i> 272:17376-17384 (1997).
,	BE	Okuley, J. <i>et al.</i> , <i>Arabidopsis FAD2</i> Gene Encodes the Enzyme That Is Essential for Polyunsaturated Lipid Synthesis. <i>The Plant Cell</i> 6:147-158 (1994).
,	BF	Polashock, J.J. <i>et al.</i> , Expression of the Yeast Δ-9 Fatty Acid Desaturase in <i>Nicotiana tabacum</i> . <i>Plant Physiol.</i> 100:894-901 (1992).
,	BG	Sayanova, O. <i>et al.</i> , Expression of a borage desaturase cDNA containing an N-terminal cytochrome <i>b_s</i> domain results in the accumulation of high levels of Δ ⁶ -desaturated fatty acids in transgenic tobacco. <i>Proc. Natl. Acad. Sci. USA</i> 94:4211-4216 (1997).
,	BH	Schultz, D.J. <i>et al.</i> , Expression of a Δ ⁹ 14:0-acyl carrier protein fatty acid desaturase gene is necessary for the production of a ω ⁵ anacardic acids found in pest-resistant geranium (<i>Pelargonium xhortorum</i>). <i>Proc. Natl. Acad. Sci. USA</i> 93:8771-8775 (1996).
,	BI	Shah, S. <i>et al.</i> , Overexpression of the <i>FAD3</i> Desaturase Gene in a Mutant of <i>Arabidopsis</i> . <i>Plant Physiol.</i> 114:1533-1539 (1997).
,	BJ	Shanklin, J. <i>et al.</i> , Eight Histidine Residues Are Catalytically Essential in a Membrane-Associated Iron Enzyme, Stearyl-CoA Desaturase, and Are Conserved in Alkane Hydroxylase and Xylene Monoxygenase. <i>Biochemistry</i> 33:12787-12794 (1994).
RK	BK	Shanklin, J. <i>et al.</i> , Mössbauer studies of alkane ω-hydroxylase: Evidence for a diiron cluster in an integral-membrane enzyme. <i>Proc. Natl. Acad. Sci. USA</i> 94:2981-2986 (1997).

EXAMINER

Ronelle Kapoor

DATE CONSIDERED

1-8-03



Sheet 3 of 3

RECEIVED
Sheet 3 of 13
JUN 21 2002
1
TECH CENTER 1600/25
Hitchcock

EXAMINER Ronell Kallis

DATE CONSIDERED

1 - 8-03